

Date:

Name:

Venous Insufficiency

CEAP Revised Classification: **C**linical, **E**tiologic, **A**natomic, **P**athophysiologic

| | | | |
|---|---------------------------------------------------|------|--|
| C | Asymptomatic | C0 | |
| | Teleangiectasia, Reticular Veins, Malleolar Flare | C1 | |
| | Varicoce veins | C2 | |
| | Edema, no skin changes | C3 | |
| | Pigmentation ,eczema | C4a | |
| | Lipodermatosclerosis, atrophic blanche | C4b | |
| | Skin change as above healed ulcer | C5 | |
| | Skin change as above active ulcer | C6 | |
| S | C level Symptomatic | S | |
| A | C level Asymptomatic | A | |
| E | Congenital | Ec | |
| | Primary | Ep | |
| | Secondary, known cause (post-thrombotic) | Es | |
| | No venous cause identified | An | |
| A | Superficial Veins (As), Telangiectasia/reticular | As1 | |
| | GSV, AK | As2 | |
| | GSV, BK | As3 | |
| | SSV | As4 | |
| | Non-Saphenous | As5 | |
| | Deep Veins (Ad) IVC | Ad6 | |
| | Common Iliac | Ad7 | |
| | Internal Iliac | Ad8 | |
| | External Iliac | Ad9 | |
| | Pelvic, Gonadal, Broad Ligament | Ad10 | |
| | Common Femoral | Ad11 | |
| | Deep Femoral | Ad12 | |
| | Femoral | Ad13 | |
| | Popliteal | Ad14 | |
| | Tibial, Posterior Tibial, Peroneal | Ad15 | |
| | Muscular, GI, Soleal | Ad16 | |
| | Perforating Veins (Ap) Thigh | Ap17 | |
| | Calf | Ap18 | |
| | No venous location identified | An | |
| P | Pathophysiology (P) Reflux | Pr | |
| | Obstruction | Po | |
| | Reflux and Obstruction | Pr,o | |
| | No venous pathophysiology identified | Pn | |
| L | Level of Investigation (L) Office | LI | |
| | Vascular Lab | LII | |
| | Venography, CTV, MRV, Venous Pressure | LIII | |

Philip L. Rice, M.D., F.A.C.S., R.P.V.I.
University Cardiothoracic and Vascular Associates
100 Radnor Road
State College, PA 16801
Telephone (814) 238-2616